

AD-A104 322

RECOGNITION OF COCCIDIODES IMMITIS ANTIGENS WITH
MONOCLONAL ANTIBODIES(U) CALIFORNIA UNIV OAKLAND NAVAL
BIOSCIENCES LAB S J KRAEGER ET AL. 30 SEP 86

1/1


UNCLASSIFIED

UC-NBL-963 N00014-81-C-0570

F/G 6/3

NL

END
9-87
DTIC



SECURITY CLASS

AD-A184 322

DOCUMENTATION PAGE

1a. REPORT SEC UNCLASSIFIED		1b. RESTRICTIVE MARKINGS NONE	
2a. SECURITY CLASSIFICATION AUTHORITY N/A		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited	
2b. DECLASSIFICATION/DOWNGRADING N/A		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NBL No. 963		7a. NAME OF MONITORING ORGANIZATION Office of Naval Research	
6a. NAME OF PERFORMING ORGANIZATION University of California	6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code) Code 1141 800 North Quincy Ave Arlington, VA 22217-5000	
6c. ADDRESS (City, State, and ZIP Code) Naval Biosciences Laboratory Naval Supply Center Oakland, California 94625	8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-81-C-0570	
8b. OFFICE SYMBOL (if applicable)	10. SOURCE OF FUNDING NUMBERS		
8c. ADDRESS (City, State, and ZIP Code) 800 North Quincy Avenue Arlington, Va 22217-5000	PROGRAM ELEMENT NO 61153N	PROJECT NO RR041-05	TASK NO RR041-05-03 WORK UNIT ACCESSION NO NR204-123
11. TITLE (Include Security Classification) (U) RECOGNITION OF COCCIDIOIDES IMMITIS ANTIGENS WITH MONOCLONAL ANTIBODIES			
12. PERSONAL AUTHOR(S) Kraeger, Sping J. Gennevois, Danial J.P., Cox, R. A., and Karu, Alexander E..			
13a. TYPE OF REPORT Summary Report	13b. TIME COVERED FROM 851001 TO 860930	14. DATE OF REPORT (Year, Month, Day) 1986	15. PAGE COUNT 2
16. SUPPLEMENTARY NOTATION In press. 1st International Symposium of Fungal Antigens, France, Sept 1986.			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
06	03	Keywords: Monoclonal antibodies, Immunogens	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) This note summarizes recent observations on the antigenic specificity and suitability for diagnostic use of seven IgM monoclonal antibodies (MAbs) prepared in 1984 with c. immitis Silveira spherules and endospore/ shperule culture filtrate (ECSF) as immunogens. Hybridoma cell line SPL. IVA5A1 was deposited at the American Type Culture Collection on Mar 29, 1984, and grated accession no. HB 8535.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Head, Biological Sciences Div, ONR		22b. TELEPHONE (Include Area Code) (202) 696-4986	
		22c. OFFICE SYMBOL ONR Code 1141	

963
Kraeger, S.J., Gennevois, D.J.P., and Karu, A.E. 1986. Recognition of *Coccidioides immitis* antigens with monoclonal antibodies. in: Proceedings of the 1st International Symposium on Fungal Antigens, Paris France, Sept. 1986. (in press)

RECOGNITION OF COCCIDIOIDES IMMITIS ANTIGENS WITH MONOCLONAL ANTIBODIES

S.J. Kraeger¹, D.J.P. Gennevois², R.A. Cox³, and A.E. Karu²

¹University of California, School of Public Health, Berkeley, CA; ²Naval Biosciences Laboratory, Oakland, CA; ³San Antonio Chest Hospital, San Antonio, TX

This note summarizes recent observations on the antigenic specificity and suitability for diagnostic use of seven IgM monoclonal antibodies (MAbs) prepared in 1984 with *C. immitis* Silveira spherules and endospore/spherule culture filtrate (ESCF) as immunogens.

Nature of the Antigens. All seven MAbs reacted with a heat-stable methanol-precipitable extract of *C. immitis* endospores, suggesting that the antigenic determinants include carbohydrate. Denaturing electrophoresis and immunoblotting of ESCF resolved protein-containing bands with apparent molecular weights of 66, 65, 56, 45, and 32 kDa that were recognized by MAbs E35 and S82, suggesting shared epitopes. Coccidioidomycosis patient sera reacted with these same bands. The same bands bound fluorescein-labeled concanavalin A (conA), but no other lectins. Competition between conA and E35 suggested that glucose or mannose is close to, or part of the epitope of E35. Non-denaturing gel filtration resolved the antigens in ESCF into a broad distribution of sizes which showed three separate patterns of reactivity with the different MAbs (Figure 1). Deproteinized extracts of the largest carbohydrate-containing antigens (fractions 21-38 and 42-43 in Figure 1) were too complex for magnetic resonance analysis, but fractions 55 and 56, which bound only MAbs E35 and S82, contained a single oligosaccharide of B-D-glucose.

Specificity of the Antibodies. Only two MAbs showed more than 20% cross-reactivity with the other major systemic fungi in radio- and enzyme immunoassays (RIA, EIA). All of the MAbs reacted with spherules of six different *C. immitis* strains and ESCF from the Silveira strain, but four MAbs did not react with the ESCF from strain 46, which is of low virulence. Fluorescence microscopy showed specific differential staining of spherules, hyphae, and arthroconidia by the MAbs. For example, Figure 2 shows staining of the ends of arthroconidia by MAb E36. The MAbs also bound differently to these morphologic forms in RIAs.

Diagnostic Applications. All seven MAbs reacted with the *C. immitis* skin-testing antigens and the mycelial lysates that are currently used to detect precipitin-forming and complement-fixing antibodies. Experiments are under way to determine whether any of the antigens are diagnostic for coccidioidomycosis. We have developed a latex agglutination (LA) test to detect *C. immitis* antigens; to date the detection limit is 1 ng of *C. immitis* ESCF at 20 ng/ml in buffer. Detection thresholds for other pathogenic fungi are

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail. and/or Special
A-1	



summarized in Table 1. We are attempting to optimize sensitivity and specificity when sera are tested by this method.

In summary, the determinants recognized by all seven MABs appear to be primarily carbohydrates, found on several biochemically separable species. Each MAB reacted differently with various antigens or fungal particles in EIA, RIA, immunofluorescence, and LA tests. MABs in this panel detect taxonomic and developmental differences in isolates of *C. immitis* and other pathogenic fungi, and some can be used to localize antigens on the surface of fungal cells. Some of these MABs are uniquely specific for *C. immitis* antigens, and their potential usefulness in diagnostic assays for coccidioidomycosis deserves further study.

Acknowledgements. We thank Dr. O. Hindsgaul, University of Alberta, for performing proton and carbon magnetic resonance analyses, and Dr. D. Baker, Bioresponse Inc., for valuable discussions. [supported by ONR Contract N00014-81-C-0570 (SJK, DG, AEK), and NIAID Grant A121431 (RAC)]

copy of
figures not
attached.

Fig. 1. Sephacryl S-400 gel filtration profile of *C. immitis* ESCF. The top panel shows the protein and carbohydrate content of each fraction. The succeeding panels show the reactivities of three MABs with equivalent amounts of carbohydrate from each column fraction in an EIA. The carbohydrate profile is reproduced in each panel for reference. V_0 is the void volume.

Fig. 2. Fluorescence micrograph of *C. immitis* arthroconidia reacted with MAB E36 and fluorescein-labeled goat anti-mouse globulin, showing differential staining of the arthroconidia. Magnification X 594.

Table 1. Minimum detectable dilutions of culture filtrates (ng of carbohydrate) in a microplate LA test. Polystyrene beads (0.3 μ m) coated with the indicated MABs were mixed with culture filtrates diluted in 0.05ml of 0.85% NaCl-0.05 M glycine-NaOH (pH 8.2) with 0.1% bovine serum albumin, in microtiter "U"-shaped wells. Agglutination was scored after incubating the plates overnight at 22 C.

END

9-87

DTIC